

Adventure Tourism

[English](#)

[Vietnamese](#)

There has been a growing interest in adventure tourism lately. However, organizing adventure tours is not an easy task. It requires very careful preparation with attention to specific details.

This tour has p young male and q female participants. In addition to the logistic and rescue team, the organizers also assign k more guides to join the tour. In the first stage of the tour, the road is quite narrow passing a cliff; the group will have to go in one line. To be able to help each other, a female participant has to go next to, i.e. before or after, a male participant or a guide.

Furthermore, there must be at least one participant next to a guide. Given these constraints, there are several ways the group can form a line. Let's denote B, G and M as a male participant, a female participant and a guide respectively. A line formation can be represented by a string of length $(p+q+k)$ containing characters from the set (B, G, M). Two line formations are different if their string representations are different. For example, the group having 2 male, 2 female and a guide ($p = q = 2, k = 1$) has 24 different way to form a line as follows:

Given $p, q,$ and $k,$ let's denote n as the number of different ways to form a line. Your task is to write a program to calculate the remainder of n divided by 10^7 .

Input

The input file consists of several data sets. The first line of the input file contains the number of data sets which is a positive integer and is not bigger than 20. The following lines describe the data sets.

For each data test, there is only one line containing three integers p, q and k ($0 \leq p, q \leq 1\,000, 0 \leq k \leq 10$) separated by space.

Output

For each data test, write in one line the remainder of the number of different line formations divided by 10^7 .

Example

Sample Input

```
1
2 2 1
```

Sample Output

```
24
```