

Sub array Sum1

Let $A = \{a_0, a_1, a_2, a_3, \dots, a_{n-1}\}$ be an array. We define a recursive operation Op on array A as follows

$$Op(A) = Op(\text{two}(A)) + Op(\text{one}(A)) + Op(\text{zero}(A)) \text{ if } n > 1 \\ = A \text{ otherwise}$$

Here, $\text{zero}(A) = \{a_0, a_3, a_6, \dots\}$ i.e. an array formed by elements whose indices are divisible by 3. Similarly, $\text{one}(A) = \{a_1, a_4, a_7, a_{10}, \dots\}$ and $\text{two}(A) = \{a_2, a_5, a_8, a_{11}, \dots\}$. Also, $+$ is the concatenation operation.

For example, if $A = \{0, 1, 2, 3, 4, 5\}$. Then $Op(A)$ will be calculated as

$$Op(A) = Op(\{2, 5\}) + Op(\{1, 4\}) + Op(\{0, 3\}) \\ = Op(\{\}) + Op(\{5\}) + Op(\{2\}) + Op(\{\}) + Op(\{4\}) + Op(\{1\}) + Op(\{\}) + Op(\{3\}) + Op(\{0\}) \\ = \{5, 2, 4, 1, 3, 0\}$$

We define an query on an array B as taking the sum of all elements b_k where $i \leq k \leq j$ and $l \leq b_k \leq r$.

Input

First line contains size n of array C . ($n \leq 10^5$) -

Second line contains n integers $c_0, c_1, c_2, \dots, c_{n-1}$. ($|c_i| < 10^6$) -

Third line contains q , number of queries. ($q \leq 10^5$) -

Next q lines contains four integers i, j, l, r . ($0 \leq i < n, i \leq j < n, l = -10^6 - 1, r = 10^6 + 1$)

Note that l, r are fixed

Output

You have to output q integers corresponding to each query on a separate line.

Example

Input:

```
4
1 -1 5 4
1
0 3 -1000001 1000001
```

Output:

```
9
```